

Datasheet for ABIN3136877

## CHMP1A Protein (AA 1-196) (His tag)



[Go to Product page](#)

### 1 Image

#### Overview

Quantity:	1 mg
Target:	CHMP1A
Protein Characteristics:	AA 1-196
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CHMP1A protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)

#### Product Details

Sequence: MDDTLFQLKF TAKQLEKLAK KAEKDSKAEQ AKVKKALQKQ NVECARVYAE NAIRKKNEGV  
NWLRMASRVD AVASKVQTAV TMKGVTKNMA QVTKALDKAL SAMDLQKVSA VMDRFEQQVQ  
NLDVHTSVME DSVSSATLTL TPQEQVDSL I VQIAEENGL E VLDQLSQLPE GASAVGESSV  
RSQEDQLSRR LAALRN

**Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.**

- Characteristics:
- Made in Germany - from design to production - by highly experienced protein experts.
  - Mouse Chmp1a Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.
  - State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

## Product Details

---

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

---

Purification:	Two step purification of proteins expressed in bacterial culture: <ol style="list-style-type: none"><li>1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li><li>2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li></ol>
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Endotoxin has not been removed. Please contact us if you require endotoxin removal.
Grade:	Crystallography grade

---

## Target Details

---

Target:	CHMP1A
Alternative Name:	Chmp1a ( <a href="#">CHMP1A Products</a> )
Background:	Probable peripherally associated component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are

---

## Target Details

---

generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I,-II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. Involved in cytokinesis. Involved in recruiting VPS4A and/or VPS4B to the midbody of dividing cells. May also be involved in chromosome condensation. Targets the Polycomb group (PcG) protein BMI1/PCGF4 to regions of condensed chromatin. May play a role in stable cell cycle progression and in PcG gene silencing (By similarity). {ECO:0000250}.

---

Molecular Weight: 22.6 kDa Including tag.

---

UniProt: [Q921W0](#)

## Application Details

---

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: Protein has not been tested for activity yet. In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

---

Expiry Date: Unlimited (if stored properly)



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process